

A 3D ribbon diagram of a protein complex, likely a transcription factor or nuclear receptor, rendered in purple, yellow, and green. The structure is complex and multi-domain, with various alpha-helices and beta-sheets visible.

Products for Transcription Biology

**Transcription Factor &
Nuclear Receptor ELISAs**

Chromatin Assays & IP Kits

**Antibodies, Cell Extracts &
Recombinant Proteins**

Luciferase Assays


**Supershift &
Gelshift**

ACTIVE  MOTIF®

Tools to Analyze
Nuclear Function



Active Motif Offices



U.S.

1914 Palomar Oaks Way, Suite 150
Carlsbad, CA 92008 USA
Toll Free: 1 877 222 9543
Direct: 1 760 431 1263
Fax: 1 760 431 1351
Email: sales@activemotif.com



EUROPE

104 Avenue Franklin Roosevelt – Box 25
B-1330 Rixensart, Belgium
Germany Free Phone: 0800/181 99 10
France Free Phone: 0800/90 99 79
UK Free Phone: 0800/169 31 47
Other countries, Direct: +32 (0)2 653 0001
Fax: +32 (0)2 653 0050
Email: eurotech@activemotif.com

JAPAN

Azuma Bldg. 7th Floor
2-21 Ageba-Cho, Shinjuku-Ku
Tokyo, 162-0824, Japan
Direct: +81 (0)3 5225 3638
Fax: +81 (0)3 5261 8733
Email: japantech@activemotif.com

Transcription Factor DNA-binding ELISAs

TransAM™ AML/Runx2
 TransAM™ AP-1.....2
 TransAM™ ATF-2.....2
 TransAM™ c-Myc.....2
 TransAM™ C/EBP α/β2
 TransAM™ CREB and pCREB2
 TransAM™ Elk-1.....2
 TransAM™ ER.....2
 TransAM™ FKHR (FOXO1).....2
 TransAM™ GATA.....2
 TransAM™ GR.....2
 TransAM™ HIF-1.....2
 TransAM™ HNF.....2
 TransAM™ IRF-3.....2
 TransAM™ IRF-7.....2
 TransAM™ MAPK.....2
 TransAM™ MEF2.....2
 TransAM™ MyoD.....2
 TransAM™ NF-YA.....2
 TransAM™ NFAT.....2
 TransAM™ NF κ B.....2
 TransAM™ NF κ B Chemi.....2
 TransAM™ NF κ B Flexi.....2
 TransAM™ Nrf2.....2
 TransAM™ Oct-4.....2
 TransAM™ p53.....2
 TransAM™ PPAR.....2
 TransAM™ Sp1 and Sp3.....2
 TransAM™ SREBP1.....2
 TransAM™ STAT.....2

Co-Immunoprecipitation

Universal Magnetic Co-IP Kit6

Chromatin Immunoprecipitation

ChIP-IT™ Express & ChIP-IT™ Express HT7
 ChIP-IT™ Protein G Magnetic Beads.....7
 ChIP-IT™ Control Kits.....7
 Ready-to-ChIP Chromatin.....7
 Re-ChIP-IT™8
 Bridging Antibody for Mouse IgG9

Luciferase Reporter Assay

RapidReporter®10

Recombinant Proteins

Recombinant Transcription Factors12
 FunctionELISA™ I κ B α12

Cell Extracts

High-quality Cell Extracts13

Antibodies

Transcription Factor and Related Antibodies14

Nuclear & Cytoplasmic Extraction

Nuclear Extract Kit16

Supershift & Gelshift Assays

Nushift™ and Gelshift™ Kits17

Nuclear Receptor Binding Assays

NR Sandwich AR.....19
 NR Peptide ER α19
 NR Sandwich ER α19
 NR Sandwich PR.....19

DISTRIBUTORS

Distribution Network.....20

On the Cover

Crystal structure of NFAT1 (green) and Fos-Jun (yellow) on the IL-2 ARRE1 (A-T-rich purine-box/antigen receptor response element) (purple) binding site.

TransAM™

TransAM™ Kits are highly sensitive ELISA-based assays* that facilitate the study of transcription factor activation in mammalian tissue and cell extracts. They combine a fast, user-friendly format with a sensitive, specific assay. The standard TransAM Kits are 10-fold more sensitive than conventional gelshift techniques (Figure 2), while TransAM Chemi Kits utilize chemiluminescent detection to improve assay sensitivity to only 40 ng of extract. The TransAM method eliminates the use of radioactivity and provides quantitative results in less than five hours, while its high-throughput format offers the flexibility to screen up to 96 samples simultaneously.

The TransAM™ advantage

TransAM Kits offer a novel method for rapid and sensitive quantitative measurement of transcription factor activation. TransAM Kits are available in the original format that contains a 96-well plate to which an immobilized oligonucleotide containing a consensus-binding site has been immobilized or in a Flexi format that enables you to bind any oligonucleotide to the plate. In both formats, cell extract is added to each well and the transcription factor of interest binds specifically to the oligonucleotide that is immobilized on the plate. Each well is then incubated with primary antibody specific for the active form of the bound transcription factor. Subsequent incubation with an HRP-conjugated

sensitive, non-radioactive transcription factor ELISAs

Until now, transcription factors have been studied using primarily three methods: gelshift assays, immunoblotting and reporter gene assays. These methods are time consuming and provide, at best, only semi-quantitative results. Moreover, they don't support high-throughput methods and lack sensitivity, specificity and reproducibility. The TransAM technology eliminates these shortcomings, making transcription factor research faster and more precise.

* TransAM is licensed from EAT under issued and pending worldwide patents. Purchase includes the right to use for basic research only. Other-use licenses available, please contact Technical Services.

secondary antibody and either the standard or chemiluminescent Developing Solution provides an easily quantified readout (Figure 1).


TransAM Kits are available for study of a variety of individual transcription factors and transcription factor families. TransAM eliminates the need for time consuming cloning, cell transfections, and gel exposures. The use of messy radioactive probes is also unnecessary. There is also no need for constructing stable reporter cell lines, or for dealing with the variability in expression of transfected reporter genes.

WHY USE TRANSAM™?

- Up to 100-fold greater sensitivity than gelshift assays
- Results in less than 5 hours
- Non-radioactive, colorimetric readout easily quantified by spectrophotometry
- No cloning or cell transfections required
- Ability to assay tissue samples
- 96-stripwell format is compatible with high-throughput automation

FIGURE 1:
Flow chart of the TransAM process.

Activated transcription factor in cell extract binds to oligonucleotides immobilized in the well. Incubation with primary and secondary antibodies quantifies the amount of activated transcription factor in a sample.



Increased sensitivity & accuracy

Small changes in transcription factor levels can have a significant impact on cellular function. Therefore, it is important to use a sensitive assay when studying transcription factor activation. TransAM Kits are 10-fold more sensitive than gelshift assays and also yield more quantitative results. To illustrate, NFκB p50 activity was assayed using the TransAM NFκB p50 Kit and EMSA (Figure 2). These data clearly demonstrates that the DNA binding ELISA format of the TransAM Kits yields more sensitive and quantitative results than traditional assays such as EMSA.

In addition, TransAM Kits provide more accurate measurements than traditional assays because of their broad linear range of detection. This eliminates the need to run multiple assays to obtain accurate results. Both the increased sensitivity and accuracy of the TransAM Kits will drastically improve your transcription factor analysis. The combination of a fast, user-friendly format with high sensitivity and specificity is validated by over 900 journal citations (Figure 4).

Proven specificity

To accurately study transcription factor activation, it is vital to be able to determine which isoform of the transcription factor of interest is involved in pathway regulation. All TransAM Kits are tested for specificity by performing the assays in the presence of an excess of oligonucleotide containing a wild-type or mutated consensus-binding site (Figure 3). This competitive assay proves that the transcription factor detected is binding specifically to the probe that has been immobilized to the TransAM plate. All TransAM antibodies are also assayed for cross-reactivity with other closely related family members to ensure that only the isoform of interest is detected.

Applications of TransAM™ Kits

TransAM Kits have unsurpassed flexibility, which is why they have been used successfully with both primary and transformed cell lines as well as tissue samples. Common applications include:

- Drug screening and/or potency studies
- Monitoring of transcriptional regulation activity
- Protein structure/function studies






FIGURE 4:
TransAM citations.
 Using HighWire Press, <http://highwire.stanford.edu>, a comparison was made by searching for citations of TransAM from Active Motif versus the tradenames of all competitor kits combined. TransAM is clearly the leader.




A




B

FIGURE 2:
TransAM Kits are more sensitive than gelshift.
 Human fibroblast WI-38 cells are stimulated with IL-1α for 30 minutes. Increasing amounts of whole-cell extract are assayed using the TransAM NFκB p50 Kit (A) or gel retardation (B). The TransAM method is 10-fold more sensitive and provides more quantitative data.



A



B

FIGURE 3:
Specificity of TransAM Kits.
 TransAM NFκB p50 Kit assays are performed in the presence of wild-type and mutated competitor oligonucleotides using 10 µg/well whole-cell extract from human fibroblast WI-38 cells stimulated with IL-1α for 30 minutes (A) and 5 µg/well whole-cell extract from HeLa cells stimulated with TNF-α for 10 minutes, 30 minutes and 2 hours (B).

TransAM™ Ordering Information

FACTOR	PRODUCT	FORMAT	Cat. No.
AML/Runx	TransAM™ AML-1/Runx1	1 x 96-well plate	47396
		5 x 96-well plates	47896
	TransAM™ AML-3/Runx2	1 x 96-well plate	44496
		5 x 96-well plates	44996
AP-1	TransAM™ AP-1 Family (c-Fos, FosB, Fra-1, Fra-2, c-Jun, JunB & JunD)	2 x 96-well plates	44296
	TransAM™ AP-1 c-Fos	1 x 96-well plate	44096
		5 x 96-well plates	44596
	Recombinant c-Fos protein	5 µg	31115
	TransAM™ AP-1 c-Jun	1 x 96-well plate	46096
		5 x 96-well plates	46596
	Recombinant c-Jun protein	5 µg	31116
	TransAM™ AP-1 FosB	1 x 96-well plate	45096
	5 x 96-well plates	45596	
	TransAM™ AP-1 JunD	1 x 96-well plate	43496
		5 x 96-well plates	43996
ATF-2	TransAM™ ATF-2	1 x 96-well plate	42396
		5 x 96-well plates	42896
c-Myc	TransAM™ c-Myc	1 x 96-well plate	43396
		5 x 96-well plates	43896
	Recombinant c-Myc protein	5 µg	31117
C/EBP	TransAM™ C/EBP α/β	1 x 96-well plate	44196
		5 x 96-well plates	44696
CREB	TransAM™ CREB	1 x 96-well plate	42096
		5 x 96-well plates	42596
	TransAM™ pCREB	1 x 96-well plate	43096
		5 x 96-well plates	43596
	Recombinant CREB protein	25 µg	31107
Elk-1	TransAM™ Elk-1	1 x 96-well plate	44396
		5 x 96-well plates	44896
ER	TransAM™ ER	1 x 96-well plate	41396
		5 x 96-well plates	41996
FKHR	TransAM™ FKHR (FOXO1)	1 x 96-well plate	46396
		5 x 96-well plates	46896
GATA	TransAM™ GATA Family (GATA-1, GATA-2 & GATA-3)	2 x 96-well plates	48296
	TransAM™ GATA-4	1 x 96-well plate	46496
		5 x 96-well plates	46996
GR	TransAM™ GR	1 x 96-well plate	45496
		5 x 96-well plates	45996
HIF	TransAM™ HIF-1	1 x 96-well plate	47096
		5 x 96-well plates	47596
HNF	TransAM™ HNF Family (HNF-1, HNF-3α, HNF-3β & HNF-4α)	2 x 96-well plates	46296
	TransAM™ HNF-1	1 x 96-well plate	46196
		5 x 96-well plates	46696
IRF-3	TransAM™ IRF-3 (Human)	1 x 96-well plate	48396
		5 x 96-well plates	48896
	TransAM™ IRF-3 (Mouse)	1 x 96-well plate	48496
		5 x 96-well plates	48996
IRF-7	TransAM™ IRF-7	1 x 96-well plate	50196
		5 x 96-well plates	50696

TransAM™ Ordering Information, continued

FACTOR	PRODUCT	FORMAT	Cat. No.	
MAPK	TransAM™ MAPK Family (ATF-2, c-Jun, c-Myc, MEF2 & STAT1α)	2 x 96-well plates	47296	
MEF2	TransAM™ MEF2	1 x 96-well plate 5 x 96-well plates	43196 43696	
MyoD	TransAM™ MyoD	1 x 96-well plate 5 x 96-well plates	47196 47696	
NF-YA	TransAM™ NF-YA	1 x 96-well plate 5 x 96-well plates	40396 40896	
NFAT	TransAM™ NFATc1	1 x 96-well plate 5 x 96-well plates	40296 40796	
NFκB	TransAM™ NFκB Family (p50, p52, p65, c-Rel & RelB)	2 x 96-well plates	43296	
	TransAM™ Flexi NFκB Family (p50, p52, p65, c-Rel & RelB)	2 x 96-well plates	43298	
	TransAM™ NFκB p50	1 x 96-well plate 5 x 96-well plates	41096 41596	
	TransAM™ NFκB p50 Chemi	1 x 96-well plate 5 x 96-well plates	41097 41597	
	TransAM™ Flexi NFκB p50	1 x 96-well plate	41098	
	Recombinant NFκB p50 protein	5 μg	31101	
	TransAM™ NFκB p52	1 x 96-well plate 5 x 96-well plates	48196 48696	
	TransAM™ NFκB p52 Chemi	1 x 96-well plate 5 x 96-well plates	48197 48697	
	TransAM™ NFκB p65	1 x 96-well plate 5 x 96-well plates	40096 40596	
	TransAM™ NFκB p65 Chemi	1 x 96-well plate 5 x 96-well plates	40097 40597	
	TransAM™ Flexi NFκB p65	1 x 96-well plate	40098	
	Recombinant NFκB p65 protein	5 μg	31102	
	Nrf2	TransAM™ Nrf2	1 x 96-well plate 5 x 96-well plates	50296 50796
	Oct-4	TransAM™ Oct-4	1 x 96-well plate 5 x 96-well plates	42496 42996
p53	TransAM™ p53	1 x 96-well plate 5 x 96-well plates	41196 41696	
	Recombinant p53 protein	5 μg	31103	
	PPAR	TransAM™ PPARγ	1 x 96-well plate 5 x 96-well plates	40196 40696
Sp1 and Sp3	TransAM™ Sp1	1 x 96-well plate 5 x 96-well plates	41296 41796	
	TransAM™ Sp1/Sp3	1 x 96-well plate 5 x 96-well plates	40496 40996	
	Recombinant Sp1 protein	5 μg	31136	
	SREBP1	TransAM™ SREBP1	1 x 96-well plate 5 x 96-well plates	50496 50996
STAT	TransAM™ STAT Family (STAT1α, STAT3, STAT5A & STAT5B)	2 x 96-well plates	42296	
	TransAM™ STAT3	1 x 96-well plate 5 x 96-well plates	45196 45696	
	Nuclear Extract Kit (see page 16 for details)	100 rxns 400 rxns	40010 40410	
	ProStain™ Protein Quantification Kit (visit www.activemotif.com for details)	1000 rxns	15001	

Universal Magnetic Co-IP Kit

lower background; suitable for both nuclear & whole-cell complexes

The Universal Magnetic Co-IP Kit improves co-immunoprecipitation (Co-IP) through the use of protein G-coated magnetic beads, which speed and simplify the IP and wash steps while greatly reducing background. The kit includes optimized

reagents for making both nuclear & whole-cell extracts from cells or tissue. This gives you the flexibility to Co-IP any protein complex, whether it was originally bound to DNA (Figure 1) or in the cytoplasm.

Co-IP cytoplasmic AND nuclear complexes

Co-IP is often used to study cytoplasmic protein complexes. But, traditional methods are not optimal for studying DNA-binding proteins because nuclear complexes are very fragile, causing them to be disrupted during extraction. For this reason, in addition to containing components for preparing whole-cell extracts, the Universal Magnetic Co-IP Kit provides nuclear extraction reagents that have been optimized to preserve nuclear protein complexes. The kit's Enzymatic Shearing Cocktail uses DNA digestion to gently release the nuclear protein complexes from the DNA, so they are intact and ready for Co-IP.

UNIVERSAL MAGNETIC CO-IP KIT ADVANTAGES

- Magnetic beads simplify procedure and reduce background
- Optimized extraction method maintains nuclear protein complexes
- Preserve protein modifications

Simpler procedure, lower background

The Universal Magnetic Co-IP Kit utilizes protein G-coated magnetic beads, which simplify Co-IP by enabling the IP and wash steps to be performed in seconds, rather than having to use centrifugation. Because these beads have very low non-specific binding, background is reduced even while using the kit's low-salt Co-IP/Wash Buffer, which helps maintain weaker complexes.

Complete kit for better results

The Universal Magnetic Co-IP Kit has both nuclear and whole-cell extraction reagents, so you can perform IP on all types of protein complexes. The kit also includes protein G-coated magnetic beads, a unique Co-IP/Wash Buffer as well as phosphatase, protease and deacetylase inhibitors that preserve the integrity of the proteins and protein modifications (Figures 2 & 3). Finally, the kit includes a strong bar magnet, so you can take advantage of the improved wash and IP steps enabled by the magnetic beads. This makes the Universal Magnetic Co-IP Kit a simple, flexible and complete solution for getting more from your Co-IP.




FIGURE 1:
Nuclear Co-IP of SRC-1 and ER α .

The Universal Magnetic Co-IP Kit was used to make nuclear extract from MCF-7 cells induced 1 hour with 10 nM Estradiol. IP was performed on 300 μ g samples using 2 μ g SRC-1 pAb, ER α pAb and rabbit IgG (as a negative control). Western blot was then performed using the ER α pAb on 10 μ g Input Extract (Lane 1), SRC-1 IP (Lane 2), ER α IP (Lane 3) and the rabbit IgG IP (Lane 4).




FIGURE 2:
Detection of acetylated Histone H3. HeLa nuclear extracts were made using the Universal Magnetic Co-IP Kit and a traditional high-salt extraction protocol, each supplemented with 1 μ M trichostatin A, a deacetylase inhibitor. Five and ten μ g samples of each extract were used in Western blot with Histone H3 acetyl rabbit pAb (Cat. No. 39139). The pan acetyl-H3 (arrow) was detected only in samples made using the kit's gentle nuclear extraction procedure.




FIGURE 3:
Preserve acetylation and methylation. Nuclear extracts were made from HeLa cells treated with 500 nM trichostatin A for 20 hours using either the Universal Magnetic Co-IP Kit (with its deacetylase inhibitor) or traditional high-salt extraction. Five and ten μ g samples of these extracts were used in Western blot with Histone H3 acetyl pAb (Cat. No. 39139) and Histone H3 trimethyl Lys4 pAb (Cat. No. 39159). The acetylated protein was detected only in the sample made using the kit. Methylation was slightly better maintained in the sample made using the kit.

Product	Format	Cat. No.
Universal Magnetic Co-IP Kit	25 rxns	54002

ChIP-IT™ Express & ChIP-IT Express HT

improved kits greatly reduce background

Active Motif has improved its ChIP-IT™ Express Kits by greatly reducing the background, which improves your results and enables you to use less starting material than ever before. In addition, the provided magnetic beads have made it possible to

streamline the protocol so you can get results in half the normal time with much less sample manipulation. These advances have made possible the new ChIP-IT Express HT Kit, which enables you to perform true high-throughput ChIP in 96-well plates.

ChIP-IT EXPRESS ADVANTAGES

- No more need for pre-clearing, blocking or DNA purification steps
- Reduced background
- High throughput compatible
- Dramatically reduced hands-on time

The most efficient ChIP enrichment kit

ChIP is an enrichment technique, not a purification method. Thus, the less efficient your enrichment, the higher the sample background and the more material you will need to obtain an interpretable result. Conventional ChIP requires at least 2 million cells as starting material, which can be problematic with some cell lines. At the least, growing this many cells is labor intensive. Active Motif's improved ChIP-IT Express Kits, however, have been optimized to provide superior target gene enrichment, resulting in unmatched sensitivity. Using ChIP-IT Express, it is routine to perform ChIP on material from as few as 750,000 cells and the kits have even been shown to work with as few as 12,500 cells (Figure 1).

High-throughput ChIP

The efficient ChIP-IT Express method has led to the recent development of ChIP-IT Express HT, which enables you to perform ChIP in 96-well plates using multi-channel pipettors.




FIGURE 1:

ChIP-IT Express works with 12,500 cells.

ChIP-IT Express was performed in duplicate on decreasing amounts of sonicated HeLa cell chromatin. Two μg of RNA pol II and Neg IgG antibody was used for IP. GAPDH PCR primers were used to analyze the immunoprecipitated DNA. Using the improved ChIP-IT Express reagents and protocol, positive ChIP data was obtained from as few as 12,500 cells.

The magnetic bead advantage

The ChIP-IT Express magnetic beads have less background than standard agarose beads, so pre-clearing and blocking steps are no longer necessary. The magnetic pull-down occurs in just seconds, and re-formulated buffers allow steps to be combined and DNA purification to be eliminated. ChIP-IT Express is available in both sonication and enzymatic shearing formats.

Positive controls ensure success

Because interpreting ChIP data can be difficult, Active Motif has developed a complete set of controls to help you understand your results and troubleshoot your assays. To provide you with controls that are appropriate for your research, we offer our human, mouse and rat ChIP-IT Control Kits separately from ChIP-IT Express Kits. These provide positive and negative control antibodies and species-specific primers, PCR buffer and a convenient 10X DNA loading dye so your PCR reactions are gel-ready. All reagents are quality control tested and validated to ensure your ChIP assay is working properly. In addition, we also offer convenient Ready-to-ChIP Chromatin from a number of ENCODE cell lines, so you can be certain that the only variable in validating a new antibody for ChIP is the antibody itself.

Try the best ChIP kits today

For additional information on the ChIP-IT Express and ChIP-IT Express HT Kits, please visit us at www.activemotif.com/chipit.

Product	Format	Cat. No.
ChIP-IT™ Express	25 rxns	53008
ChIP-IT™ Express Enzymatic	25 rxns	53009
ChIP-IT™ Express HT	96 rxns	53018
ChIP-IT™ Protein G Magnetic Beads	25 rxns	53014
ChIP-IT™ Control Kit – Human	5 rxns	53010
ChIP-IT™ Control Kit – Mouse	5 rxns	53011
ChIP-IT™ Control Kit – Rat	5 rxns	53012
Ready-to-ChIP HeLa Chromatin	10 rxns	53015
Ready-to-ChIP Hep G2 Chromatin	10 rxns	53019
Ready-to-ChIP K-562 Chromatin	10 rxns	53020
Ready-to-ChIP NIH/3T3 Chromatin	10 rxns	53021

Re-ChIP-IT™

identify protein co-localization *in vivo* using sequential chromatin IP

Performing sequential chromatin IP (also called Re-ChIP) was technically challenging and difficult, until now. Active Motif's Re-ChIP-IT™ Kit makes it easy to perform sequential ChIP, so you

can localize two different proteins or histone modifications to the same genomic locus.

Extend the utility of ChIP

When performing chromatin immunoprecipitation (ChIP) experiments, it is often useful to prove that two different proteins or histone modifications are present at the same site in the genome. Or, you may want to determine if a protein coincides with a specific histone modification at the same regulatory element. Re-ChIP (aka Sequential ChIP, Chromatin Re-IP and ChIP Re-ChIP) is a relatively new technique in which sequential chromatin immunoprecipitations are performed using two different antibodies. This enables you to assay for the simultaneous presence of two proteins or distinct histone modifications at the same genomic region of interest.

Active Motif's new Re-ChIP-IT Kit makes it simple for you to perform these types of experiments. All buffers for chromatin IP are included, making it easy to get started. And, the detailed protocol ensures you're successful the first and every time. Plus, Active Motif offers a variety of ChIP controls kits (page 9) to help you validate the results of your Re-ChIP experiments.




FIGURE 2: Schematic representation of the Re-ChIP-IT procedure.




FIGURE 1: Sequential chromatin immunoprecipitation using Re-ChIP-IT. The lane numbers are the same in each panel to indicate that the DNA is from the same chromatin sample. The left panel shows the results of PCR performed on an aliquot of DNA removed from the experiment after the first ChIP step; the right panel represents PCR results on DNA from chromatin samples after both ChIP steps. For example, chromatin samples subjected to first ChIP using Mouse IgG as a negative control (lanes 1 and 2 in the left panel) were then subjected to a second ChIP with an RNA Pol II antibody (lanes 1 and 2 in the right panel). Chromatin samples in which Mouse IgG was used as either the first antibody (lanes 1 and 2) or second antibody (lanes 5 and 6) show little amplification of GAPDH DNA in either the left (first ChIP) or right panel (first and second ChIP). Chromatin samples in which the first antibody used was anti-RNA Pol II and the second antibody was anti-TFIIB (lanes 3 and 4) show good amplification of GAPDH DNA after the second ChIP (right panel) indicating co-localization of RNA Pol II and TFIIB at the same region of the GAPDH promoter.

Sequential chromatin IP made easy

Re-ChIP-IT uses magnetic beads that have less background than standard agarose beads, so pre-clearing and blocking steps are not needed. Magnetic pull-down occurs in just seconds, and the method's low background has eliminated the need for DNA purification. And, Re-ChIP-IT can be used with chromatin prepared using our sonication or enzymatic shearing kits.

Get started with Re-ChIP

For additional information on the new Re-ChIP-IT Kit go to www.activemotif.com/rechip.

Product	Format	Cat. No.
Re-ChIP-IT™	25 rxns	53016

ChIP Accessory Kits and Reagents

maximize your chromatin IP experiments

ChIP & IP enhancer for mouse antibodies

Low antibody binding affinity can make getting good ChIP & IP results difficult. Because mouse IgM and IgG₁ antibodies do not have strong binding affinity for protein G, ChIP and regular immunoprecipitation (IP) can be challenging when using protein G-conjugated beads. The Bridging Antibody for Mouse IgG facilitates improved binding of mouse antibodies, especially IgG₁ and IgM isotypes, to protein G to generate more robust binding and better results in ChIP and standard IP experiments (Figure 1).




FIGURE 1:
Improvement in chromatin IP using an anti-mouse bridging antibody.
 ChIP was performed using chromatin from U-937 cells induced with TNF- α . PCR was performed with primers corresponding to the human IL-8 promoter.
 Lanes 1-4: Beads not pre-incubated with bridging antibody.
 Lanes 5-8: Beads pre-incubated with 5 μ g bridging antibody.
 Lanes 1, 2 & 5, 6: ChIP performed using anti-p65 mouse mAb, 2 μ g per IP.
 Lanes 3, 4 & 7, 8: ChIP performed using negative control mouse IgG.

Beads, controls and shearing kits

Active Motif offers a broad range of reagents and accessory kits to complement the ChIP-IT™ line of ChIP kits. These products will help you prepare chromatin, troubleshoot your ChIP experiments and make them more reproducible.




FIGURE 2:
Analysis of DNA sheared using the Enzymatic Shearing Kit.
 HeLa cells were fixed for 10 minutes with 1% formaldehyde and then chromatin was prepared using the Enzymatic Shearing Kit protocol. Chromatin was sheared with the Enzymatic Shearing Cocktail for 5, 10 or 15 minutes and the reaction was stopped.
 Lane 1: 100 to 1000 bp ladder.
 Lane 2: Unsheread HeLa DNA.
 Lane 3: HeLa DNA treated for 5 minutes.
 Lane 4: HeLa DNA treated for 10 minutes (optimized).
 Lane 5: HeLa DNA treated for 15 minutes.

Ready-to-ChIP Chromatin

For your convenience, Active Motif offers Ready-to-ChIP Chromatin from a number of ENCODE cell lines, which have been optimally sheared by sonication and validated in ChIP. As a result, you can more easily validate your own antibodies and primer sets. The chromatin can be used with all of the ChIP-IT Kits and controls, so you can be certain that the only variable in validating a new antibody for ChIP is the antibody itself.

Product	Format	Cat. No.
Bridging Antibody for Mouse IgG	500 μ g	53017

Product	Format	Cat. No.
ChIP-IT™ Protein G Magnetic Beads	25 rxns	53014
ChIP-IT™ Control Kit – Human	5 rxns	53010
ChIP-IT™ Control Kit – Mouse	5 rxns	53011
ChIP-IT™ Control Kit – Rat	5 rxns	53012
Ready-to-ChIP HeLa Chromatin	10 rxns	53015
Ready-to-ChIP Hep G2 Chromatin	10 rxns	53019
Ready-to-ChIP K-562 Chromatin	10 rxns	53020
Ready-to-ChIP NIH/3T3 Chromatin	10 rxns	53021
ChIP-IT™ Shearing Kit	10 rxns	53002
Enzymatic Shearing Kit	10 rxns	53005

RapidReporter®

double destabilization improves luciferase assay response and sensitivity

Reporter gene assays are an important tool used for studying signal transduction pathways because of their simplicity and versatility. However, standard reporter assays are limited by the fact that the basal activity of the cloned promoter results in accumulation of luciferase mRNA and protein. The slow clearance rate of these reporter molecules substantially delays and dilutes the measurable response to stimulation or repression. As a result, with standard reporter assays, transient or relatively minor effects are hidden and kinetic assays are inaccurate. To solve this problem, some assays incorporate a protein-destabilizing element in the vector, causing the luciferase to degrade more rapidly. However, this only partly addresses the problem as reporter clearance rates are also dependent on the half-life of the reporter mRNA.

Active Motif's patented RapidReporter®* method overcomes these limitations by utilizing vectors that include both protein and mRNA destabilizing elements. This causes the cell to rapidly degrade both the luciferase protein and mRNA, which lowers background and increases both the magnitude of the response that can be measured after stimulation or repression, and the speed with which the assay can report changes in transcription. In addition, RapidReporter uses a modified *Gussia* luciferase, the most sensitive luciferase known. The combination of *Gussia* expression and double destabilization means that RapidReporter enables researchers to detect smaller changes in activity than is possible using non-destabilized reporter gene assays, or those that destabilize the protein only.

The RapidReporter® advantage

RapidReporter vectors utilize a non-secreted form of the extremely bright *Gussia* luciferase gene, and are available as "empty" vectors with a multiple cloning site (MCS) for insertion of promoters/enhancers, or as pre-made vectors that contain a widely studied promoter such as NFκB, CREB (CRE) or GR. In the RapidReporter method, cells are plated and transiently transfected with the appropriate RapidReporter vector. After an overnight incubation, the stimulator/repressor of transcriptional activity is added for the appropriate time. The cells are lysed and flash luminescence is measured. All vectors are available separately or as a complete assay kit, which includes an EFlα promoter-driven positive control vector, as well as Lysis and Assay Buffers and a substrate optimized for use with the *Gussia* luciferase, which guarantees the best results possible.

WHY USE RAPIDREPORTER®?

- Stronger fold induction reduces false positives
- Faster response enables detection of transient effects, and drugs that decompose rapidly
- Your choice of stringency – pRR-High is highly destabilized for best response to changes in transcription, while pRR-Low contains fewer destabilizing elements for a stronger basal signal
- Improved performance increases productivity and lead identification
- Offered as "empty" vectors for cloning your own elements or as pre-mades, which contain widely studied promoter sites





FIGURE 1: RapidReporter provides a better fold induction and more accurate kinetics.

293 cells were transiently transfected with pRR-High-CRE, pRR-Low-CRE and pGL3 vector containing CRE (CREB response element) and plated onto 96-well plates. Twenty-four hours post-transfection, cells were stimulated with 4 μM isoproterenol. At the indicated time points, the cells were measured for *Gussia* luciferase (pRR) and firefly luciferase activities (pGL3).

Your choice of stringency

RapidReporter vectors are offered in two different stringencies, pRR-High and pRR-Low. The pRR-High vector provides the lowest background possible because it contains more and stronger destabilization elements, thereby providing maximum responsiveness. The fold induction of stimulated vs. non-stimulated samples is highest using pRR-High whereas the basal (background) level will be lowest. This makes pRR-High vectors ideal in cases where your stimulation will only produce a weak effect. In contrast, the pRR-Low vector contains fewer and weaker destabilization elements. While this lowers the level of fold-induction observed, it enables weaker basal signals to be detected. pRR-Low vectors are ideal for use when a low signal strength is expected, such as with cells that transfect only at a low efficiency or when testing a promoter with only weak activity in the cells of interest. Regardless of the RapidReporter vector used, the fold induction observed will be higher than with other luciferase reporter systems such as firefly (Figure 1) and *Renilla* (data not shown).



Pre-made reporter vectors

The RapidReporter vectors were designed with the *Gaussia* luciferase reporter and an ampicillin resistance gene for plasmid selection. The multiple cloning site offers maximum flexibility for insertion, whatever cloning method is used. For your convenience, Active Motif also offers a number of pre-made vectors that already contain widely studied promoters. Each pre-made is available in the pRR-High vector (Figure 2) to maximize responsiveness to changes in transcription.

Optimized buffers and substrates

Active Motif's *Gaussia* Luciferase Assay Kit contains proprietary Lysis and Assay Buffers and a *Gaussia* Substrate, which are optimized for use with the *Gaussia* luciferase-fusion proteins expressed by the RapidReporter vectors. This enables you to take advantage of the enhanced sensitivity of pRR-High and pRR-Low plasmids. Because the *Gaussia* luciferase gene encoded by pRR-High and -Low vectors has been modified for intracellular expression, it is not suitable for use with standard *Gaussia* assay kits or assay kits designed for use with other luciferases. For your convenience the RapidReporter *Gaussia* Luciferase Assay Kits are available in both 100 and 1000 reaction formats.

FIGURE 2:

The RapidReporter pRR-High vector.

The pRR-High plasmid contains two strong protein-destabilizing elements (PDE), while pRR-Low contains only one PDE. Each vector also contains an RNA-destabilizing element (RDE); the RDE in the pRR-High vector is strong while the RDE in the pRR-Low vector is weaker. The vectors also contain an intron for improved expression by enhancing RNA processing, and an optimized synthetic poly A signal on the reporter gene to enhance its expression.

Product	Format	Cat. No.
RapidReporter® <i>Gaussia</i> Luciferase Assay	100 rxns	33001
	1000 rxns	33002
RapidReporter® pRR-High vector	10 µg	33003
RapidReporter® pRR-High Assay	100 rxns	33004
RapidReporter® pRR-Low vector	10 µg	33005
RapidReporter® pRR-Low Assay	100 rxns	33006
RapidReporter® pRR-High-CRE vector	10 µg	33007
RapidReporter® pRR-High-CRE Assay	100 rxns	33008
RapidReporter® pRR-High-GR vector	10 µg	33011
RapidReporter® pRR-High-GR Assay	100 rxns	33012
RapidReporter® pRR-High-IRF-1 vector	10 µg	33017
RapidReporter® pRR-High-IRF-1 Assay	100 rxns	33018
RapidReporter® pRR-High-NFκB vector	10 µg	33009
RapidReporter® pRR-High-NFκB Assay	100 rxns	33010
RapidReporter® pRR-High-STAT1 vector	10 µg	33015
RapidReporter® pRR-High-STAT1 Assay	100 rxns	33016
RapidReporter® pRR-High-STAT3 vector	10 µg	33013
RapidReporter® pRR-High-STAT3 Assay	100 rxns	33014

* RapidReporter is covered under U.S. Patent No. 7,157,272 and various other patents worldwide, is a registered trademark of, and is sold under license granted by GeneStream Pty Ltd. Purchasers are subject to a Limited-use Label License that allows them to use RapidReporter for a 6-month evaluation period. Commercial entities and all users who wish to perform high-throughput screening will be required to enter into an End User License Agreement with GeneStream following the evaluation period. Please contact Active Motif's Technical Services for details.

Recombinant Proteins

purified proteins for all your research

Active Motif provides an extensive line of recombinant proteins, including transcription factors and cell signaling-related proteins. These recombinant proteins are ideal for all your research needs and can be used in a variety of applications including as protein standards in ELISAs. The c-Fos, c-Jun, c-Myc, CREB, NFκB p50, NFκB p65, p53 and Sp1 proteins have been validated for use in

making standard curves in our TransAM™ Transcription Factor ELISAs (see page 2). The list below is only a few of the over 100 proteins available; please visit www.activemotif.com/proteins for a complete product listing plus information on the recombinant proteins, including detailed technical data sheets that specify the protein length, the method used for purification, etc.

PRODUCT	EXPRESSED IN	FORMAT	Cat. No.
Recombinant c-Fos protein	Baculovirus	5 µg	31115
Recombinant c-Jun protein	Baculovirus	5 µg	31116
Recombinant c-Myc protein	<i>E. coli</i>	5 µg	31117
Recombinant CREB protein	<i>E. coli</i>	25 µg	31107
Recombinant eIF2α protein	Sf9 cells	25 µg	31108
Recombinant NFκB p50 protein	<i>E. coli</i>	5 µg	31101
Recombinant NFκB p65 protein	<i>E. coli</i>	5 µg	31102
Recombinant p53 protein	<i>E. coli</i>	5 µg	31103
Recombinant p300 protein	Baculovirus	4 µg	31124
Purified Sp1 protein	HeLa cells	2 µg	31137
Recombinant STAT1 protein	<i>E. coli</i>	25 µg	31112
Recombinant STAT3 protein	<i>E. coli</i>	10 µg	31140

FunctionELISA™ IκBα

study the phosphorylation state of IκBα

NFκB is regulated by the IκB family of inhibitory proteins. Phosphorylation of IκBα leads, ultimately, to activation of NFκB. Thus, analysis of the phosphorylation state of IκBα provides insights about NFκB and the many genes it regulates (see page 2 for information on TransAM™ NFκB Kits). FunctionELISA IκBα uses

a Capture Antibody specific for phosphorylated IκBα. It utilizes highly sensitive chemiluminescent detection to enable accurate study of the phosphorylation state of IκBα (Figure 1). Function-ELISA IκBα requires the use of a luminometer.

Applications

FunctionELISA™ IκBα Kits can be used to study the phosphorylation state of IκBα and to correlate this information with the activation and translocation of NFκB. Applications include drug screening and/or potency effects on phosphorylation of IκBα and time course experiments to monitor the duration of IκBα phosphorylation.

Product	Format	Cat. No.
FunctionELISA IκBα	1 x 96-well plate	48005
	5 x 96-well plates	48505




FIGURE 1:

Induction of IκBα phosphorylation.

Jurkat cells were treated with 1 nM TNF-α and harvested at the indicated time points. Whole-cell extracts were assayed in Western blot analysis (A) using IκBα phospho Ser32,36 mAb (Cat. No. 40904) and the FunctionELISA IκBα Kit (B).

Cell Extracts

high-quality extracts ensure reproducible results

Active Motif provides high-quality extracts from a variety of cell types and tissue sources. Our extracts are available in nuclear, cytoplasmic and whole-cell formats, using extraction protocols that have been optimized to produce high yields of cell signaling and transcription factor-related proteins. In addition, many of our extracts are prepared from cells that have been stimulated or treated to specifically induce activation of hard-to-detect transcription factors.

Active Motif's ready-to-use extracts are ideal for use as positive controls in a variety of applications, including TransAM™ Kits (see pages 2-5), gelshift and supershift assays, Western blot analysis or as a starting point for transcription factor purification. As the list below is only a small subset of the over 100 extracts available, please visit www.activemotif.com/extracts for a complete product listing and more complete information.

EXTRACT	FORMAT	Cat. No.
3T3 nuclear extract	200 µg	36000
3T3 nuclear extract (1 hr serum response)	200 µg	36001
3T3-L1 nuclear extract	200 µg	36071
COS-7 nuclear extract	200 µg	36079
COS-7 nuclear extract (CoCl ₂ treated)	200 µg	40600
COS-7 nuclear extract (IFN γ treated)	200 µg	36112
HeLa whole-cell extract	200 µg	40050
HeLa nuclear extract	200 µg	36010
HeLa whole-cell extract (IL-1 α stimulated)	200 µg	40100
HeLa whole-cell extract (TNF- α stimulated)	200 µg	40200
HeLa nuclear extract (2 hr serum response)	200 µg	36104
HeLa nuclear extract (4 hr serum response)	200 µg	36086
HeLa nuclear extract (Anisomycin treated)	200 µg	36111
HeLa nuclear extract (TNF- α stimulated)	200 µg	40210
HeLa nuclear extract (TPA stimulated, 10 minutes)	200 µg	36109
HeLa nuclear extract (TPA stimulated, 2 hours)	200 µg	36009
Hep G2 cytoplasmic extract	200 µg	36400
Hep G2 nuclear extract	200 µg	36011
Hep G2 nuclear extract (Acetaldehyde treated)	200 µg	36065
Hep G2 nuclear extract (IFN γ treated)	200 µg	36093
Hep G2 whole-cell extract (IL-6 stimulated, 10 ng/ml)	200 µg	36413
Hep G2 nuclear extract (IL-6 stimulated, 10 ng/ml)	200 µg	36092
Hep G2 nuclear extract (IL-6 stimulated, 100 ng/ml)	200 µg	36107
HL-60 nuclear extract	200 µg	36072
Jurkat cytoplasmic extract	200 µg	36402
Jurkat nuclear extract	200 µg	36014
Jurkat nuclear extract (Heat Shock)	200 µg	36069

EXTRACT	FORMAT	Cat. No.
Jurkat nuclear extract (IFN γ treated)	200 µg	36094
Jurkat nuclear extract (PHA stimulated)	200 µg	36108
Jurkat cytoplasmic extract (TPA + CI stimulated)	200 µg	36401
Jurkat nuclear extract (1 day growth)	200 µg	36110
Jurkat nuclear extract (TPA + CI stimulated)	200 µg	36013
MCF-7 nuclear extract	200 µg	36017
MCF-7 nuclear extract (Estradiol treated)	200 µg	36016
MCF-7 nuclear extract (H ₂ O ₂ -treated)	200 µg	40800
MCF-7 nuclear extract (H ₂ O ₂ post-treated)	200 µg	40810
MCF-7 nuclear extract (Spermine treated)	200 µg	36068
Nb2 nuclear extract	200 µg	36105
Nb2 nuclear extract (prolactin stimulated)	200 µg	36106
NIH/3T3 cytoplasmic extract	200 µg	36403
NIH/3T3 nuclear extract	200 µg	36020
NIH/3T3 nuclear extract (IL-6 stimulated)	200 µg	36056
NIH/3T3 whole-cell extract (TGF- β 1 treated)	200 µg	36414
NIH/3T3 whole-cell extract (TPA stimulated)	200 µg	36415
NIH/3T3 nuclear extract (1 hr serum response)	200 µg	36021
U-937 cytoplasmic extract	200 µg	36410
U-937 nuclear extract	200 µg	36030
U-937 nuclear extract (IFN α treated)	200 µg	36077
U-937 cytoplasmic extract (TPA stimulated)	200 µg	36411
U-937 nuclear extract (TPA stimulated)	200 µg	36029
WI-38 nuclear extract	200 µg	40310
WI-38 nuclear extract (Forskolin stimulated)	200 µg	40300
WI-38 nuclear extract (TPA + CI stimulated)	200 µg	40500

Antibodies

highly characterized to meet your needs

Active Motif offers a broad line of antibodies directed against a variety of transcription factor, histone and cell signaling-related proteins. These antibodies are ideal for performing CHIP, Western blots, ELISAs, immunoprecipitation and supershift/electrophoretic mobility shift assays (EMSA), as indicated in the table below. To ensure their high quality, all antibodies are characterized using

appropriate cellular extracts and the specificity of each antibody is verified by peptide competition. We also offer a large range of cell extracts for use as positive controls, or in your functional studies (see page 13). The list below is only a portion of the 100's of antibodies available; please visit www.activemotif.com/abs for a complete product listing and more complete information.

ANTIBODY	IMMUNOGEN	REACTIVITY	APPLICATIONS	FORMAT	Cat. No.
AML-1/Runx1 pAb	Raised against amino acids 231-245 of human AML-1/Runx1.	H, M	WB	100 µl	39000
AML-2/Runx3 pAb	Raised against full-length human AML-2/Runx3.	H	SS, WB	100 µl	39301
AML-3/Runx2 pAb	Raised against amino acids 333-348 of human AML-3/Runx2.	H	SS	100 µl	39302
c-Fos pAb	Raised against amino acids 358-372 of human c-Fos.	H, M	WB	100 µl	39008
c-Fos pAb	Raised against amino acids 358-372 of human c-Fos.	H, M	SS	100 µl	39308
c-Jun pAb	Raised against amino acids 20-37 of human c-Jun.	H, M	WB	100 µl	39009
c-Jun pAb	Raised against amino acids 20-37 of human c-Jun.	H, M	CHIP, IF, SS	100 µg	39309
c-Myc mAb (61 kDa form)	Raised against amino acids 408-420 of human c-Myc; detects 61 kDa form.	H	WB	100 µl	39502
c-Myc pAb (61 kDa form)	Raised against amino acids 9-22 of human c-Myc; detects 65 kDa form.	H, M	WB	100 µl	39011
c-Myc pAb (65 kDa form)	Raised against amino acids 9-22 of human c-Myc.	H, M, MK	WB	100 µl	39012
c-Rel pAb	Raised against amino acids 507-524 of human c-Rel.	H	SS, WB	100 µg	39311
C/EBPα pAb	Raised against amino acids 1-14 of human C/EBPα.	H, R	CHIP, IF, SS, WB	100 µg	39306
C/EBPβ pAb	Raised against amino acids 228-242 of human C/EBPβ.	H, R	SS, WB	100 µl	39307
CREB mAb	Raised against a synthetic peptide to human CREB.	H	WB	100 µg	40961
CREB-1 pAb	Raised against amino acids 6-23 of human CREB-1.	H, R	WB	100 µl	39013
CREB-1 pAb	Raised against amino acids 6-23 of human CREB-1.	H, R	SS	17 rxns	39310
E2F-1 mAb	Raised against full-length human E2F-1.	H	SS	17 rxns	39601
E2F-1 pAb	Raised against amino acids 309-323 of human E2F-1.	H, M	CHIP, SS	17 rxns	39313
E2F-3 mAb	Raised against full-length human E2F-3.	H	WB	100 µl	39507
E2F-3 mAb	Raised against full-length human E2F-3.	H	SS	17 rxns	39602
E2F-4 mAb	Raised against full-length human E2F-4.	H	WB	100 µl	39508
E2F-4 mAb	Raised against full-length human E2F-4.	H	SS	17 rxns	39603
E2F-6 mAb	Raised against full-length human E2F-6.	H	CHIP, WB	100 µl	39509
ERα mAb	Raised against full-length human ERα.	H	WB	100 µl	39510
ERα pAb	Raised against amino acids 145-159 of human ERα.	H	WB	100 µl	39021
ERα pAb	Raised against amino acids 145-159 of human ERα.	H, M	SS	17 rxns	39316
FosB mAb	Raised against amino acids 2-14 of human FosB.	H	WB	100 µg	40960
FosB pAb	Raised against amino acids 2-14 of human FosB.	H, M	WB	100 µl	39022
FosB pAb	Raised against amino acids 2-14 of human FosB.	H, M	SS	17 rxns	39318
FosB pAb	Raised against amino acids 2-14 of human FosB.	H	WB	100 µg	40959
Fra-2 pAb	Raised against amino acids 289-305 of human Fra-2.	H, M	WB	100 µl	39023
HDAC1 pAb	Raised against amino acids 1-5, 433-448 & 467-482 of hHDAC1.	H	WB	100 µg	40967
HDAC3 pAb	Raised against amino acids 2-17 of human HDAC3.	H	CHIP, WB	100 µg	40968
HDAC4 pAb	Raised against amino acids 194-209 of human HDAC4.	H	CHIP, WB	100 µg	40969
HDAC5 pAb	Raised against amino acids 572-587 of human HDAC5.	H, M	CHIP, WB	100 µg	40970
HDAC6 pAb	Raised against amino acids 1-16 of human HDAC6.	H, M	CHIP, WB	100 µg	40971
HDAC11 pAb	Raised against amino acids 238-251 of human HDAC11.	H, M	WB	200 µl	39208

REACTIVITY KEY: B = Bovine; H = Human; M = Mouse; MK = Monkey; R = Rat

ANTIBODY	IMMUNOGEN	REACTIVITY	APPLICATIONS	FORMAT	Cat. No.
HNF-1 α pAb	Raised against amino acids 540-555 of human HNF-1 α .	H, M, R	WB	100 μ l	39030
HNF-1 α pAb	Raised against amino acids 540-555 of human HNF-1 α .	H, M, R	SS	17 rxns	39321
HNF-3 α pAb	Raised against amino acids 150-165 of human HNF-3 α .	H, R	SS	17 rxns	39322
HNF-3 β pAb	Raised against amino acids 295-311 of human HNF-3 β .	H	WB	100 μ l	39096
HNF-3 γ pAb	Raised against amino acids 287-299 of human HNF-3 γ .	H, R	WB	100 μ l	39031
HNF-3 γ pAb	Raised against amino acids 287-299 of human HNF-3 γ .	H	SS	17 rxns	39323
I κ B α phospho Ser 32, 36 mAb	Raised against a peptide with phospho Ser32 & Ser36 on human I κ B α .	H	WB	100 μ g	40904
IKK α mAb	Raised against full-length human IKK α .	H, M	IP, WB	100 μ g	40905
IKK β mAb (Clone 10AG2)	Raised against full-length human IKK β .	H	WB	100 μ g	40906
IKK β mAb (Clone 10A9B6)	Raised against full-length human IKK β .	H	IP, WB	100 μ g	40907
IKK γ mAb	Raised against full-length human IKK γ .	H	WB	100 μ g	40908
IRF-1 pAb	Raised against amino acids 126-140 of human IRF-1.	H	SS	17 rxns	39325
IRF-3 pAb	Raised against full-length human IRF-3.	B, H	ChIP, WB	100 μ l	39033
IRF-6 pAb	Raised against amino acids 358-372 of human IRF-6.	H	WB	100 μ g	39080
IRF-7 pAb	Raised against amino acids 463-477 of human IRF-7.	H	WB	100 μ g	39081
JunB pAb	Raised against amino acids 3-17 of human JunB.	H, M	WB	100 μ l	39034
JunB pAb	Raised against amino acids 3-17 of human JunB.	H	ChIP, SS	17 rxns	39326
JunD pAb	Raised against amino acids 231-245 of human JunD.	H	WB	100 μ g	39082
JunD pAb	Raised against amino acids 143-157 of human JunD.	H	ChIP, SS	100 μ l	39328
NAK1/Nur77 pAb	Raised against amino acids 251-266 of human NAK1/Nur77.	H, M	WB	200 μ l	40982
NF-1A pAb	Raised against amino acids 478-492 of human NF-1A.	H, R	WB	100 μ l	39036
NF-1A pAb	Raised against amino acids 478-492 of human NF-1A.	H, R	SS	100 μ l	39329
NF κ B p50 pAb	Raised against amino acids 363-376 of human NF κ B p50.	H, M	WB	100 μ l	39037
NF κ B p50 pAb	Raised against amino acids 363-376 of human NF κ B p50.	H, M, R	SS	17 rxns	39330
NF κ B p65 pAb	Raised against amino acids 2-17 of human NF κ B p65.	H, M, R	WB	100 μ l	39038
NF κ B p65 pAb	Raised against amino acids 501-515 of human NF κ B p65.	H, R	WB	100 μ l	39039
NF κ B p65 mAb	Raised against amino acids 526-539 of human NF κ B p65.	H, M	FC, WB	100 μ g	40916
Oct-2 mAb	Raised against full-length human Oct-2.	H	WB	100 μ l	39517
Oct-2 mAb	Raised against full-length human Oct-2.	H	SS	17 rxns	39604
p53 pAb	Raised against amino acids 367-381 of human p53.	H, M	WB	100 μ l	39041
p53 pAb	Raised against amino acids 367-381 of human p53.	H	ChIP, SS	17 rxns	39334
Pax-5 pAb	Raised against amino acids 1-15 of human Pax-5.	H	WB	100 μ l	39043
Pax-5 pAb	Raised against amino acids 1-15 of human Pax-5.	H	SS	17 rxns	39336
RelB pAb	Raised against amino acids 563-579 of human RelB.	H	WB	100 μ l	39053
RelB pAb	Raised against amino acids 476-490 of human RelB.	H	SS	17 rxns	39339
RNA pol II mAb	Raised against a synthetic peptide YSPTSpPS of human RNA pol II.	H, M, R	ChIP, ELISA, IF, IP, WB	200 μ l	39097
Sp1 pAb	Raised against full-length human Sp1.	H	WB	100 μ l	39057
Sp1 pAb	Raised against amino acids 520-534 of human Sp1.	H, M, R	WB	100 μ l	39058
Sp3 pAb	Raised against full-length human Sp3.	H, M	SS	17 rxns	39341
STAT1 α pAb	Raised against amino acids 712-750 of human STAT1 α .	H	WB	100 μ l	39059
STAT2 pAb	Raised against amino acids 830-849 of human STAT2.	H, M	WB	100 μ g	39060
STAT3 pAb	Raised against full-length human STAT3.	H	WB	100 μ l	39061
STAT4 pAb	Raised against amino acids 731-748 of mouse STAT4.	H, M, R	WB	100 μ l	39062
STAT5A pAb	Raised against full-length human STAT5A.	H	WB	100 μ l	39063
STAT5B pAb	Raised against amino acids 777-787 of human STAT5B.	H	WB	100 μ l	39064
STAT6 pAb	Raised against amino acids 808-826 of human STAT6.	H	WB	100 μ l	39065
TRF2 pAb	Raised against full-length human TRF2.	H, M, R	ChIP, IP, WB	100 μ g	39223
VDR pAb	Raised against amino acids 5-19 of human VDR.	H, R	WB	100 μ l	39069
VDR pAb	Raised against amino acids 5-19 of human VDR.	H, R	SS	17 rxns	39344
YY1 pAb	Raised against amino acids 109-123 of human YY1.	H, M, MK, R	WB	100 μ l	39071
YY1 pAb	Raised against amino acids 109-123 of human YY1.	H	SS	17 rxns	39345

APPLICATIONS KEY: ChIP = Chromatin Immunoprecipitation; FC = Flow Cytometry; IF = Immunofluorescence; IP = Immunoprecipitation; SS = Supershift; WB = Western Blot

Nuclear Extract Kit

high yields of specifically segregated extracts

The Nuclear Extract Kit is ideal for the preparation of nuclear, whole-cell and cytoplasmic extracts from mammalian cells and tissues. The resultant high-quality extracts may be used with Active Motif's TransAM™ Kits (see page 12) or in gelshift assays, Western blots, DNA footprinting or as a starting point for transcription factor purification.

The Nuclear Extract Kit eliminates the need to optimize reagents and ensures consistently high yields. The detailed protocol helps ensure that your extract is not contaminated with proteins from other cellular compartments (Figure 1).

The Nuclear Extract Kit advantage

In the Nuclear Extract Kit, cells are collected in ice-cold PBS in the presence of phosphatase inhibitors to limit further protein modifications. Next, the cells are resuspended in hypotonic buffer to swell the cell membrane. Addition of detergent causes leakage of the cytoplasmic proteins into the supernatant. After collection of the cytoplasmic fraction, the nuclei are lysed and the nuclear proteins are solubilized in lysis buffer in the presence of the protease inhibitors. Whole-cell extracts can also be prepared by collecting the cells in the PBS/phosphatase inhibitors solution and lysing in lysis buffer. Solubilized proteins are separated from the cell debris by centrifugation. The concentration of protein in the cell extract is then measured by Active Motif's ProStain Protein Quantification Kit (Cat. No. 15001) or with a Bradford-based assay.




FIGURE 1: Specific extraction of nuclear and cytoplasmic extracts. Nuclear and cytoplasmic extracts were prepared using the Nuclear Extract Kit from HeLa samples unstimulated or stimulated with TNF- α for 30 minutes and assayed using the TransAM™ NF κ B p50 Kit. Because activated NF κ B translocates to the nucleus, only nuclear extract from stimulated cells should contain activated NF κ B.

WHY USE THE NUCLEAR EXTRACT KIT?

- Quality-controlled reagents ensure reproducibility
- No need to optimize your own procedure
- Complete kit contains all required reagents
- Prepare nuclear, cytoplasmic or whole-cell extracts with one kit
- Ability to prepare extracts from both cultured cells and tissue samples

Product	Format	Cat. No.
Nuclear Extract Kit	100 rxns	40010
	400 rxns	40410

Nushift™ and Gelshift™

reliable transcription factor detection

Active Motif's Nushift™ and Gelshift™ Kits offer a convenient, flexible format for performing traditional electrophoretic mobility shift assays (EMSA). These kits enable you to screen different nuclear extracts or gene promoters for specific transcription factor:DNA binding activity. Nushift Kits are designed to provide consistent supershift results, while Gelshift Kits are ideal for performing gelshift assays; each are supplied with pre-optimized components that save you both time and money. For screening transcription factor families, Active Motif offers a wide variety of Nushift and Gelshift Family Kits (see the table below).

Nushift and Gelshift Kits come complete with optimized buffers, wild-type and mutant oligonucleotides, ready-to-label wild-type oligonucleotide probe and positive control nuclear extracts. Nushift Kits also include antibody(ies) for performing supershift assays and, if available, neutralizing peptides. For your convenience, the optimized Nushift and Gelshift buffers are also available separately (see page 18). Active Motif also offers its Nuclear Extract Kit (see page 16), which can be used to prepare nuclear, cytoplasmic or whole-cell extracts from mammalian cells or tissues. The resultant high-quality extract is ideal for EMSA studies.

Nushift™ and Gelshift™ Family Kits

PRODUCT	DESCRIPTION	REACTIVITY	Cat. No.
Nushift™ AML/Runx Family	AML/Runx and AML-3/Runx2 wild-type, mutant and ready-to-label oligonucleotides; AML-1/Runx1, AML-2/Runx3 and AML-3/Runx2 rabbit polyclonal antibodies; HL-60, Raji and Saos-2 nuclear extracts.	Human	37000
Nushift™ AP-1 Family	AP-1 wild-type, mutant and ready-to-label oligonucleotides; c-Fos, c-Jun, FosB, JunB and JunD rabbit polyclonal antibodies and peptides; 3T3 (1 hr serum response), 3T6 and Jurkat (TPA + CI) nuclear extracts.	Human, Mouse	37003
Gelshift™ AP-1 Family (Carcinoma)	AP-1 wild-type, mutant and ready-to-label oligonucleotides; HeLa and HeLa (TPA) nuclear extracts.	Human, Mouse	37301
Gelshift™ AP-1 Family (Serum Response)	AP-1 wild-type, mutant and ready-to-label oligonucleotides; 3T3 and 3T3 (1 hr serum response) nuclear extracts.	Human, Mouse	37326
Gelshift™ AP-1 Family (T Cell Activated)	AP-1 wild-type, mutant and ready-to-label oligonucleotides; Jurkat and Jurkat (TPA + CI) nuclear extracts.	Human, Mouse	37302
Gelshift™ AP-1 Family (TPA-Inducible)	AP-1 wild-type, mutant and ready-to-label oligonucleotides; HeLa (TPA) and Jurkat (TPA + CI) nuclear extracts.	Human, Mouse	37303
Nushift™ Cell Cycle Regulators	E2F wild-type, mutant and ready-to-label oligonucleotides; cyclin A, DP-1, E2F-1, E2F-3, E2F-4, p107 and Rb antibodies; Raji nuclear extract.	Human	37008
Gelshift™ Cell Cycle Regulators	E2F wild-type, mutant and ready-to-label oligonucleotides; Raji nuclear extract.	Human	37331
Gelshift™ NFκB Family (Carcinoma)	NFκB wild-type, mutant and ready-to-label oligonucleotides; HeLa and HeLa (TPA) nuclear extracts.	Human	37315
Gelshift™ NFκB Family (T Cell Activated)	NFκB wild-type, mutant and ready-to-label oligonucleotides; Jurkat and Jurkat (TPA + CI) nuclear extracts.	Human	37316
Gelshift™ NFκB Family (TPA-Inducible)	NFκB wild-type, mutant and ready-to-label oligonucleotides; HeLa (TPA) and Jurkat (TPA + CI) nuclear extracts.	Human	37333
Gelshift™ NFκB/Rel Family	NFκB wild-type, mutant and ready-to-label oligonucleotides; HeLa (TPA) and Jurkat (TPA + CI) nuclear extracts.	Human	37319

Nushift™ Kits

Nushift Kits are ideal for performing traditional supershift assays. Each kit includes antibody (17 assays), neutralizing peptides (15 assays, if available), wild-type and mutant oligonucleotides (20 assays), ready-to-label wild-type oligonucleotide probe (500 assays), positive control nuclear extract (50 assays), reaction buffers and G-25 purification columns. Please visit our website for complete details on each kit.

PRODUCT	REACTIVITY	Cat. No.
Nushift™ AML-1/Runx1	Human	37001
Nushift™ AML-2/Runx3	Human	37046
Nushift™ AML-3/Runx2	Human	37002
Nushift™ AP-2	Human	37004
Nushift™ AP-2	Human	37047
Nushift™ ATF-2	Human	37005
Nushift™ C/EBP α	Human, Rat	37006
Nushift™ C/EBP β	Human, Rat	37007
Nushift™ c-Fos	Human, Mouse	37010
Nushift™ c-Jun	Human, Mouse	37011
Nushift™ CREB-1	Human, Rat	37012
Nushift™ c-Rel	Human	37013
Nushift™ cyclin A	Human	37014
Nushift™ DP-1	Human	37048
Nushift™ E2F-1	Human	37015
Nushift™ E2F-3	Human	37049
Nushift™ E2F-4	Human, Mouse	37016
Nushift™ E47	Human, Mouse	37017
Nushift™ Egr-1	Human	37018
Nushift™ ER	Human	37019
Nushift™ FosB	Human, Mouse	37050
Nushift™ GATA-1	Human	37020
Nushift™ HNF-1 α	Human, Rat	37022
Nushift™ HNF-3 α	Human, Rat	37023
Nushift™ HNF-3 γ	Human	37051
Nushift™ HSF-1	Human, Rat	37052
Nushift™ IRF-1	Human	37024
Nushift™ JunB	Human	37053
Nushift™ NF κ B p50	Human	37026
Nushift™ NF-1A	Human, Rat	37025
Nushift™ Oct-2	Human	37054
Nushift™ p107	Human, Mouse	37056
Nushift™ p53	Human, Mouse, Rat	37029
Nushift™ Pax-3	Human, Mouse	37031
Nushift™ Pax-5	Human	37032
Nushift™ Pit-1	Human, Mouse, Rat	37033
Nushift™ PPAR γ	Human	37034
Nushift™ Rb	Human	37035
Nushift™ RelB	Human	37036
Nushift™ Sp3	Human, Mouse	37038
Nushift™ SRF	Human	37057
Nushift™ USFI	Human, Rat	37041
Nushift™ VDR	Human	37042
Nushift™ YY1	Human, Mouse, Rat	37043

Gelshift™ Kits

Gelshift Kits optimize traditional gelshift assays. Each kit includes wild-type and mutant oligonucleotides (20 assays), ready-to-label wild-type oligonucleotide probe (500 assays), positive control nuclear extract (50 assays), buffers and G-25 purification columns.

PRODUCT	REACTIVITY	Cat. No.
Gelshift™ AML-1/Runx1	Human	37300
Gelshift™ AML-2/Runx3	Human	37328
Gelshift™ AML-3/Runx2	Human	37329
Gelshift™ AP-2	Human	37304
Gelshift™ ATF-2	Human	37330
Gelshift™ C/EBP	Human, Rat	37305
Gelshift™ CREB	Human, Rat	37306
Gelshift™ c-Rel	Human	37307
Gelshift™ E12/E47	Human, Mouse	37308
Gelshift™ ER	Human	37309
Gelshift™ GATA-1	Human, Mouse	37310
Gelshift™ HNF-1	Human, Rat	37332
Gelshift™ HNF-3	Human, Rat	37311
Gelshift™ HNF-4	Human, Rat	37312
Gelshift™ IRF-1	Human	37313
Gelshift™ MyoD	Human, Mouse	37314
Gelshift™ NF κ B p50	Human	37317
Gelshift™ NF κ B p65	Human, Mouse	37318
Gelshift™ Oct-2	Human	37334
Gelshift™ Oct-4	Human, Rat	37335
Gelshift™ p107	Human, Mouse	37336
Gelshift™ Pax-3	Human, Mouse	37337
Gelshift™ Pax-5	Human	37338
Gelshift™ Pit-1	Human, Mouse, Rat	37320
Gelshift™ PPAR	Human	37321
Gelshift™ SF-1	Human	37339
Gelshift™ Spl/Sp3	Human	37322
Gelshift™ SRF	Human	37323
Gelshift™ STAT1	Human	37324
Gelshift™ Tal-1	Human	37340
Gelshift™ USFI	Human, Rat	37325

Nushift™ and Gelshift™ Buffers

To save you the time and inconvenience of optimizing your own buffers, Nushift and Gelshift buffers are also available separately.

PRODUCT	FORMAT	Cat. No.
Binding Buffer B-1	240 μ l	37480
Binding Buffer B-2	240 μ l	37481
Binding Buffer B-3	240 μ l	37482
Binding Buffer B-4	240 μ l	37483
Binding Buffer C-1	240 μ l	37484
Binding Buffer C-2	240 μ l	37485
Binding Buffer C-3	240 μ l	37486
Dilution Buffer A	240 μ l	37487
Stabilizing Solution D	360 μ l	37488
MicroSpin™ G-25 Columns	5 columns	37489

Nuclear Receptor ELISAs

two quantitative methods to study nuclear receptors

Active Motif offers two innovative kit formats for studying nuclear receptors (NR). The NR Peptide ELISA kits enable quantitative measurement of ligand-activated nuclear receptor proteins, while the NR Sandwich ELISA kits provide a quick and easy method to measure the total amount of a nuclear receptor in a given sample. Both methods combine a fast, user-friendly format in a highly sensitive ELISA-based format.

Nuclear receptors belong to a superfamily of structurally related ligand-inducible transcription factors whose members include receptors for estrogen (ER), glucocorticoid (GR), androgen (AR), thyroid hormone (TR), progesterone (PR) and vitamin D (VDR). Inappropriate signaling by these receptors is associated with numerous disease states, including cancer, asthma, infertility, osteoporosis and arthritis, making them drug targets.

WHY USE NR PEPTIDE & SANDWICH ELISAs?

- NR Peptide measures ligand-activated NR
- Quantitative results in just hours
- ELISA format eliminates the need to run, blot and develop gels
- NR Sandwich measures total NR
- Analyze multiple samples in low volume, high throughput
- Ability to assay both cellular extracts and recombinant proteins

NR Peptide ELISA method

Active Motif's NR Peptide ELISAs are specifically designed to capture ligand-activated nuclear receptor. Each NR Peptide Kit provides a 96-well plate coated with Capture Peptide that includes the consensus-binding motif of the nuclear receptor's co-activator. Addition of your sample results in binding of the activated nuclear receptor to the immobilized Capture Peptide. Each well is then incubated with a specific primary antibody. Subsequent incubation with HRP-conjugated antibody and developing solution provides an easily quantified readout.




FIGURE 1:
Reproducibility of NR Sandwich ELISAs.

Increasing amounts of nuclear extract from two different prostate cancer cell lines, LNCaP and DU-145, were assayed for levels of androgen receptor using the NR Sandwich AR Kit. Each dilution was assayed in duplicate and the experiment was repeated two times. The results confirm that LNCaP cells are positive for AR, while DU-145 cells are AR negative; it also exhibits very low inter- and intra-plate variation, which demonstrates the high reproducibility of the NR Sandwich AR Kit.

NR Sandwich ELISA method

To measure total levels of a given nuclear receptor within a sample, Active Motif's NR Sandwich Kits provide a simple, rapid and quantitative alternative to traditional Western blots, EMSA and reporter assays. The NR Sandwich ELISA kits use two antibodies that recognize different epitopes on the protein of interest. The Capture Antibody, which is bound to an ELISA plate, binds the nuclear receptor protein when the sample is added. The Detecting Antibody then binds to the captured nuclear receptor protein. Subsequent incubation with a HRP-conjugated secondary antibody provides an easily quantified readout of total nuclear receptor protein levels. Because the response of nuclear receptors to a single ligand can vary between cell types, nuclear receptor assays need to be highly reproducible. Therefore, our NR Sandwich ELISAs are rigorously tested to ensure the highest level of reproducibility (Figure 1).

Product	Format	Cat. No.
NR Peptide ER α	1 x 96-well plate	49096
	5 x 96-well plates	49596
NR Peptide ER α Chemi	1 x 96-well plate	49097
	5 x 96-well plates	49597
NR Sandwich AR	1 x 96-well plate	49196
	5 x 96-well plates	49696
NR Sandwich ER α	1 x 96-well plate	49296
	5 x 96-well plates	49796
Recombinant ER Protein	4,000 units	31119
NR Sandwich PR	1 x 96-well plate	49396
	5 x 96-well plates	49896

AUSTRALIA

Australian Biosearch

P.O. Box 86
Karrinyup, WA 6921
Australia
Tel: +61 (0)8 9344 1200
Fax: +61 (0)8 9345 1259
customerservice@aust-biosearch.com.au
www.aust-biosearch.com.au

AUSTRIA

THP Medical Products

Shuttleworthstraße 19
A-1210 Wien, Austria
Tel: +43 1 292 82 80
Tel: +43 1 292 82 80 88
r.englisch@thp.at
www.thp.at

CANADA

CEDARLANE Laboratories Ltd.

4410 Paletta Court
Burlington, Ontario, Canada
L7L 5R2
Toll Free Tel: 1 800 268 5058
Tel: 289 288 0001
Fax: 289 288 0020
general@cedarlanelabs.com
www.cedarlanelabs.com/canada/

MJS BioLynx Inc.

300 Laurier Blvd.
P.O. Bag 1150
Brockville, Ontario, Canada
K6V 5W1
Toll Free Tel: 1 888 593 5969
Tel: 613 498 2126
Fax: 613 342 1341
sales@biolynx.ca
tech@biolynx.ca
www.biolynx.ca

CHINA

TaKaRa Biomedical Technology (Beijing) Co., Ltd.

Life Science Park
22 KeXueYuan Road
Changping District
Beijing, 102206, China
Tel: +86 10 8072 0985/6/7/8
Fax: +86 10 8072 0989
service@takarabiomed.com.cn
www.takarabiomed.com.cn

DENMARK

Biotech Line A/S

Industrivej 3
Slangerup
DK-3550, Denmark
Tel: +45 7027 9920
Fax: +45 7027 9929
biotechline@biotechline.dk
www.biotechline.dk

FINLAND

Finnzymes Oy

Keilaranta 16 A
Espoo
FIN-02150, Finland
Tel: +358 9 2472 3010
Fax: +358 9 2472 3200
fz@finnzymes.fi
www.finnzymes.fi

HUNGARY

BIO-KASZTEL Ltd.

22 Kelenhegyi
1118-Budapest, Hungary
Tel 1: 36-1-3853887
Tel 2: 36-1-3810694
Fax: 36-1-3810695
info@kasztel.hu
www.kasztel.hu

INDIA

Genex Life Sciences Private Limited

A - 3 Ground Floor, Meghmala
St. Janabai Road
Vile Parle (East)
India
Tel: +91 - 9892291064 / 9870091064
Fax: +91 22 - 26127069
info@genexlife.com
www.genexlife.com

ISRAEL

Almog Diagnostic

P.O. Box 459
Shoham 73142
Israel
Tel: +972 3 9773390
Fax: +972 3 9773391
almog_ya@bezeqint.net
www.almog.co.il

ITALY

Vinci-Biochem

Via Ponte di Bagnolo, 10
I-50059 Vinci (Fi)
Italy
Tel: +39 0571 568 147
Fax: +39 0571 568 132
vb@vincibiochem.it
www.vincibiochem.it

KOREA

benebiosis Co., Ltd.

911 KGIT Sangam Center
1601 Sangam-Dong
Mapo-gu
Seoul, 121-270, Korea
Tel: 82 2 322 8250
Fax: 82 2 322 8007
info@benebiosis.com
www.benebiosis.com

NEW ZEALAND

Australian Biosearch

P.O. Box 86
Karrinyup, WA 6921
Australia
Tel: +61 (0)8 9344 1200
Fax: +61 (0)8 9345 1259
customerservice@aust-biosearch.com.au
www.aust-biosearch.com.au

PORTUGAL

Frilabo Ltd.

Lda Rua S. Roque da Lameira
Porto, 71 4350-307
Portugal
Tel 1: ++351 21 849 57 51
Tel 2: ++351 225 188 912
Fax 1: ++351 21 460 31 44
Fax 2: ++351 225 188 914
fga.frilabo@iol.pt
frilabo@iol.pt

SINGAPORE

SciMed (Asia) Pte Ltd.

1k 196 Pandan Loop #07-11
Pantech Industrial Complex
Singapore 128384
Tel: 65 6779 3388
Fax: 65 6266 3086
sales@scimed.com.sg
www.scimed.com.sg

SPAIN

CondaLab

C/La Forja, 9
28850 Torrejon de Ardoz.
Madrid, Spain
Tel: +34 91 761 02 00
Fax: +34 91 656 82 28
tech@condalab.com
www.condalab.com

SWEDEN

Fisher Scientific AB

Södra Långebergsgatan 30
Göteborg / V. Frolunda
SE-412 32, Sweden
Tel: +46 31 68 94 00
Fax: +46 31 68 07 17
info@thermofisher.com
www.fishersci.se

TAIWAN

Bertec Enterprise Co., Ltd.

6F No. 800 Jungjeng Road
Junghe City 235
Taipei, Taiwan, R.O.C.
Tel: 886 2 2228 1234
Fax: 886 2 2228 1111
bertec@bertec.com.tw
www.bertec.com.tw


TURKEY

Genmar Teshis Urunleri TIC, LTD, STI

1421 SOK. No: 39/6
Alsancak-Izmir
Turkey
Tel: 0090 232 464 5609
Fax: 0090 232 464 8667
genmar@superonline.com



Active Motif Offices



U.S.


1914 Palomar Oaks Way, Suite 150
Carlsbad, CA 92008 USA
Toll Free: 1 877 222 9543
Direct: 1 760 431 1263
Fax: 1 760 431 1351
Email: sales@activemotif.com

EUROPE

104 Avenue Franklin Roosevelt – Box 25
B-1330 Rixensart, Belgium
Germany Free Phone: 0800/181 99 10
France Free Phone: 0800/90 99 79
UK Free Phone: 0800/169 31 47
Other countries, Direct: +32 (0)2 653 0001
Fax: +32 (0)2 653 0050
Email: eurotech@activemotif.com

JAPAN

Azuma Bldg. 7th Floor
2-21 Ageba-Cho, Shinjuku-Ku
Tokyo, 162-0824, Japan
Direct: +81 (0)3 5225 3638
Fax: +81 (0)3 5261 8733
Email: japantech@activemotif.com



U.S.

1914 Palomar Oaks Way, Suite 150
Carlsbad, CA 92008 USA
Toll Free: 1 877 222 9543
Direct: 1 760 431 1263
Fax: 1 760 431 1351
Email: sales@activemotif.com

EUROPE

104 Avenue Franklin Roosevelt – Box 25
B-1330 Rixensart, Belgium
Germany Free Phone: 0800/181 99 10
France Free Phone: 0800/90 99 79
UK Free Phone: 0800/169 31 47
Other countries, Direct: +32 (0)2 653 0001
Fax: +32 (0)2 653 0050
Email: eurotech@activemotif.com

JAPAN

Azuma Bldg. 7th Floor
2-21 Ageba-Cho, Shinjuku-Ku
Tokyo, 162-0824, Japan
Direct: +81 (0)3 5225 3638
Fax: +81 (0)3 5261 8733
Email: japantech@activemotif.com

ACTIVE  MOTIF®

1914 Palomar Oaks Way, Suite 150
Carlsbad, CA 92008
USA